

INFINITESIMAL RIGIDITY OF A COMPACT HYPERBOLIC 4-ORBIFOLD WITH TOTALLY GEODESIC BOUNDARY

TARIK AOUGAB AND PETER A. STORM

ABSTRACT. Kerckhoff and Storm conjectured that compact hyperbolic n -orbifolds with totally geodesic boundary are infinitesimally rigid when $n > 3$. This paper verifies this conjecture for a specific example based on the 4-dimensional hyperbolic 120-cell.

1. INTRODUCTION

Given a discrete subgroup Γ of a semisimple Lie group G , mathematicians have long studied the question of when Γ can be deformed inside G . On one side lie the great rigidity theorems of Calabi, Weil, Mostow, and Margulis, which roughly state that lattices in most semisimple Lie groups have no deformations. On the other side lie the beautiful deformations theories of discrete groups of 2×2 matrices, famous exceptions to the rigidity theorems. If G is a Lie group that is not represented by 2×2 matrices, and Γ is not a lattice, then nearly nothing is known about the possible deformations of Γ . Do reasonable geometric conditions exist that guarantee rigidity, or flexibility?

As a first step in this direction, Kerckhoff and Storm studied deformations of a specific discrete subgroup of $\text{Isom}(\mathbb{H}^4)$, the isometry group of hyperbolic 4-space [5]. This led them to the following rigidity conjecture.

Conjecture 1.1. *If discrete group $\Gamma < \text{Isom}(\mathbb{H}^n)$ has Fuchsian ends, is convex cocompact, and $n > 3$, then the inclusion map of Γ is infinitesimally rigid.*

Qualitatively, groups Γ satisfying the hypotheses of Conjecture 1.1 are very close to lattices. Drop any of the three conditions on Γ , and the conjecture becomes false. For any n , there are many convex cocompact discrete groups in $\text{Isom}(\mathbb{H}^n)$ that are not rigid, for example if Γ is a free group. For $n < 4$ there exists a rich deformation theory applicable to discrete subgroups with Fuchsian ends. Even weakening the convex cocompact condition to geometric finiteness makes the conjecture false. A counterexample is presented in [5]. Nonetheless, it seems reasonable to expect Conjecture 1.1 to be true.

Before trying to prove the full conjecture, it seems prudent first to verify it in a nontrivial case. This is our goal here. More specifically, we will construct an explicit reflection group in $\text{Isom}(\mathbb{H}^4)$ satisfying the hypotheses of Conjecture 1.1, and verify that it is infinitesimally rigid. The reflection group will be an infinite index subgroup of the lattice generated by reflections in the 3-cells of the right-angled hyperbolic 120-cell in $\mathbb{H}^4$. This discrete group was chosen carefully to make

Date: September, 2008.

Aougab and Storm were partially supported by NSF grant DMS-0741604.

the computations as simple as possible. The group is described precisely in Sections 3 and 4.

The authors thank Daniel Allcock for his help with this research.

2. PRELIMINARIES

This section is a brisk introduction to a few necessary facts from hyperbolic geometry. For a detailed introduction see, for example, Ratcliffe's book [6].

Recall that Minkowski $(n+1)$ -space, denoted as $\mathbb{M}^{n+1}$ is a real $(n+1)$ -dimensional vector space equipped with the nondegenerate symmetric bilinear form

$$(\vec{x}, \vec{y}) := -x_0y_0 + \sum_{i=1}^n x_iy_i.$$

A vector with positive norm is a space-like vector, a vector of norm 0 is light-like, and a vector with negative norm time-like. The hyperboloid model of hyperbolic n -space $\mathbb{H}^n$ is simply the set of points

$$\{x \in \mathbb{M}^{n+1} \mid (x, x) = -1 \text{ and } x_0 > 0\},$$

with the metric induced by $\mathbb{M}^{n+1}$. Let $O(1, n)$ be the group of real $(n+1) \times (n+1)$ matrices A such that $A^T M A = M$, where M is the $(n+1) \times (n+1)$ diagonal matrix with diagonal entries $\{-1, 1, 1, \dots, 1\}$. In other words, $O(1, n)$ is the group of linear transformations preserving the bilinear form $(\cdot, \cdot)$. Then the group $\text{Isom}(\mathbb{H}^n)$ of isometries of $\mathbb{H}^n$ is the open subgroup of $O(1, n)$ given by matrices preserving the hyperboloid $\mathbb{H}^n$. Note that orientation reversing isometries are included here; $\text{Isom}(\mathbb{H}^n)$ has two connected components. Throughout the paper we will think of $\text{Isom}(\mathbb{H}^n)$ as this explicit group of matrices. To simplify notation, define $G_n := \text{Isom}(\mathbb{H}^n)$.

Consider a discrete finitely generated subgroup Γ of G_n with presentation

$$\Gamma = \langle r_1, r_2, \dots, r_M \mid w_1 = w_2 = \dots = w_N = 1 \rangle,$$

where each w_i is a word in the generators r_j . (The letter r stands for “reflection”, which will soon be the focus.) To avoid degenerate cases, let us assume that Γ is not virtually abelian, and does not preserve a copy of $\mathbb{H}^m$ for $m < n$. If Γ is torsion-free, then the quotient $\mathbb{H}^n/\Gamma$ will be a hyperbolic manifold. More generally, the quotient will be an orbifold. If the quotient space has finite volume then Γ is called a lattice. Here we will be mainly interested in infinite volume quotient spaces, where a little more terminology is required.

Definition 2.1. *The discrete group Γ has Fuchsian ends if there exists a closed Γ -invariant convex set $C_\Gamma \subseteq \mathbb{H}^n$ that is an n -manifold with nonempty totally geodesic boundary, and such that the quotient C_Γ/Γ has finite volume.*

Note that the boundary of C_Γ will not be connected. In the setting of groups with Fuchsian ends, Γ is convex cocompact if and only if the quotient C_Γ/Γ is compact.

If Γ has Fuchsian ends, then ∂C_Γ consists of an infinite number of disjoint totally geodesic hyperplanes, each isometric to $\mathbb{H}^{n-1}$. When Γ is torsion-free, it has Fuchsian ends if and only if the quotient C_Γ/Γ is a finite volume hyperbolic n -manifold with nonempty totally geodesic boundary. In general, C_Γ/Γ will be a finite volume orbifold with totally geodesic boundary. For the reader familiar with convex cores, we note that if Γ has Fuchsian ends then C_Γ is the convex core of Γ .

Consider the representation variety $\text{Hom}(\Gamma, G_n)$. The slightly larger space $\text{Hom}(\Gamma, \text{O}(1, n))$ can be represented explicitly as the zero set of a collection of real polynomials as follows. A homomorphism $\rho : \Gamma \rightarrow \text{O}(1, n)$ is determined uniquely by the M matrices $\rho(r_j)$ of $\text{O}(1, n)$, which can be thought of as a point in a real vector space V of dimension $M \cdot (n + 1)^2$. The $\rho(r_j)$ must lie in $\text{O}(1, n)$, meaning for each j the $(n + 1)^2$ polynomials $\rho(r_j)^T M \rho(r_j) = M$ are satisfied. Each relation w_i of Γ becomes a system of $(n + 1)^2$ polynomial equations in the entries of the matrices $\rho(r_j)$. The variety $\text{Hom}(\Gamma, \text{O}(1, n)) \subset V$ is precisely the set of points where these $(M + N) \cdot (n + 1)^2$ polynomial equations $\{P_\alpha\}$ are satisfied. Finally, we are interested in the subset $\text{Hom}(\Gamma, G_n) \subset \text{Hom}(\Gamma, \text{O}(1, n))$. It is the set of connected components where the upper-left matrix entry of each $\rho(r_j)$ is positive. (This entry cannot be zero for a matrix in $\text{O}(1, 4)$.)

The group G_n acts on $\text{Hom}(\Gamma, G_n)$ by conjugation as follows:

$$(g \cdot \rho)(\gamma) := g\rho(\gamma)g^{-1},$$

where $g \in G_n$ and $\gamma \in \Gamma$. It is easy to see that this action is algebraic. The inclusion map $\Gamma \rightarrow G_n$ is a point $\iota \in \text{Hom}(\Gamma, G_n)$. In general, analyzing the orbits of this G_n -action can be complicated. However, using our assumptions that Γ is not virtually abelian and does not preserve a lower dimensional hyperbolic space, it is easy to show that the orbit $G_n \cdot \iota \subseteq \text{Hom}(\Gamma, G_n)$ is a manifold of dimension equal to that of G_n [4].

A homomorphism $\rho \in \text{Hom}(\Gamma, G_n)$ is *locally rigid* if it has an open neighborhood contained inside the orbit $G_n \cdot \rho$, in other words all nearby representations are obtained by conjugation. The infinitesimal analogue of this notion is useful. An *infinitesimal deformation* of ρ is a tangent vector $p'(0) \in T_\rho V$ to a smooth path $p : (-\varepsilon, \varepsilon) \rightarrow V$ such that

$$\frac{d}{dt}\bigg|_0 P_\alpha(p(t)) = 0$$

for the $(M + N) \cdot (n + 1)^2$ polynomial equations defining $\text{Hom}(\Gamma, \text{O}(1, n))$ (and locally defining $\text{Hom}(\Gamma, G_n)$). This slightly odd definition is necessitated by the possibility that $\text{Hom}(\Gamma, G_n)$ is singular. Similarly, an *infinitesimal conjugation* of ρ is a tangent vector $p'(0) \in T_\rho V$ to a smooth path p in the orbit $G_n \cdot \rho$. We say ρ is *infinitesimally rigid* if every infinitesimal deformation is an infinitesimal conjugation. An infinitesimally rigid homomorphism is also locally rigid [9]. By the above discussion, to show that the inclusion map ι of G_n is infinitesimally rigid, it suffices to show that the linear subspace of infinitesimal deformations has dimension equal to that of G_n .

When $\Gamma < G_n$ is a lattice, we have the famous rigidity theorem of Calabi, Weil, and Garland.

Theorem 2.2. [1, 9, 3] *If $\Gamma < \text{Isom}(\mathbb{H}^n)$ is a lattice and $n > 3$ then the inclusion map of Γ is infinitesimally rigid.*

For discrete groups that are not lattices, Theorem 2.2 is false, but it is interesting to study whether or not a similar rigidity theorem might hold for any other natural class of discrete groups. Looking for infinite covolume groups which are as “close” to lattices as possible, Kerckhoff and Storm were led to consider discrete groups with Fuchsian ends. In $\mathbb{H}^3$, groups with Fuchsian ends are not rigid, and have a beautiful deformation theory investigated by many people [7]. In higher dimensions, simple naive dimension counts suggest that groups with Fuchsian ends are rigid.

These dimension counts are, of course, not rigorous. Nonetheless, Kerckhoff and Storm were led to the following conjecture, stated first in the introduction.

Conjecture 1.1. *If discrete group $\Gamma < \text{Isom}(\mathbb{H}^n)$ has Fuchsian ends, is convex cocompact, and $n > 3$, then the inclusion map of Γ is infinitesimally rigid.*

Note that Conjecture 1.1 is false without the assumption that Γ is convex cocompact. A counterexample was studied in [5]. As a first step toward proving this conjecture, the goal of this paper is to verify it in a specific 4-dimensional example. In search of an example where the computations are as simple as possible, the first place to look is among hyperbolic reflection groups, which we now introduce briefly. For more information the authors recommend [8].

A reflection isometry in $G_4 = \text{Isom}(\mathbb{H}^4)$ is given by a matrix of $O(1, 4)$ with a 4-dimensional eigenspace of eigenvalue 1, and a single eigenvalue equal to -1 whose corresponding eigenvector is space-like. It fixes a codimension 1 totally geodesic hyperplane of $\mathbb{H}^4$ given by the intersection of its $+1$ -eigenspace with $\mathbb{H}^4$. The orthogonal complement of the fixed hyperplane is reflected across the hyperplane by the isometry. The reflection isometry is determined uniquely by the hyperplane and vice versa.

Consider the interesting special case where $\Gamma < G_4$ is a group generated by M reflections $r_1, r_2, \dots, r_M$ with corresponding hyperplanes $H_1, H_2, \dots, H_M$, and the hyperplanes bound a convex (possibly infinite volume) region P of $\mathbb{H}^4$ known as a fundamental domain. Moreover, assume the dihedral angles of P are all $\pi/2$. In this case the presentation of Γ takes the following nice form [8]:

$$\Gamma = \langle r_1, r_2, \dots, r_M \mid (r_i r_j)^{m_{ij}} = r_i^2 = 1 \rangle,$$

where for each pair $1 \leq i < j \leq M$, m_{ij} is either 2 or infinity. As usual, the “relation” $(r_i r_j)^\infty = 1$ indicates the product $r_i r_j$ has infinite order. The product of reflections $r_i r_j$ has order 2 when the hyperplanes H_i and H_j intersect at angle $\pi/2$. This is best seen by thinking about the picture in the plane of the product of two reflections. Otherwise the product has infinite order, or $m_{ij} = \infty$. (Here we are not allowing intersections at other angles.)

We will be considering the representation variety $\text{Hom}(\Gamma, G_4) \subset \mathbb{R}^{25M}$ for groups Γ of the above form.

The following lemma shows that near the inclusion map in $\text{Hom}(\Gamma, G_4)$ any representation has the property that it maps the generators r_i to reflection isometries of G_4 .

Lemma 2.3. *Let $A \in G_n$ be a reflection isometry. Then A has an open neighborhood $U \subset G_n$ such that if $B \in U$ and $B^2 = \text{Id}$, then B is a reflection isometry.*

Proof: Recall that any involution can be diagonalized over $\mathbb{C}$ such that the diagonal entries are all ± 1 . Recall also that a reflection must have exactly one eigenvalue equal to -1 . We can define the open neighborhood U of A to be all matrices in G_n having $(n-1)$ positive eigenvalues (not necessarily distinct) and 1 negative eigenvalue. Then any matrix in U that is an involution will necessarily be a nontrivial isometry which fixes a hyperplane of codimension 1, in other words, a reflection. $\square$

Corollary 2.4. *There exists an open neighborhood $U \subset \text{Hom}(\Gamma, G_4) \subset \mathbb{R}^{25M}$ of the inclusion map such that if $\rho \in U$, then $\rho(r_i)$ is a reflection for all i .*

The representation variety $\text{Hom}(\Gamma, G_4)$ sits naturally as (connected components of) a subvariety of $\mathbb{R}^{25M}$, but by exploiting the fact that locally all the generators are reflections, it is possible to reduce the dimension considerably. To a reflection isometry r with fixed hyperplane H we can associate the 1-dimensional subspace of $\mathbb{M}^5$ given by vectors Minkowski orthogonal to H . This subspace is simply the $+1$ -eigenspace of the matrix corresponding to r , and it will consist of space-like vectors. In reverse, choosing a space-like vector $\vec{n}$ determines a codimension 1 hyperplane $\vec{n}^\perp \cap \mathbb{H}^4$, which in turn determines a reflection isometry. Letting $\mathcal{S} \subset \mathbb{M}^5$ denote the set of space-like vectors, this process defines a map

$$\mu : \mathcal{S}^M \rightarrow G_4^M \subset \mathbb{R}^{25M}$$

with image equal to the set of M -tuples of reflection isometries.

Moreover, the geometry of the fundamental domain P can be read from the normal vectors. Specifically, let $\vec{n}_1$ and $\vec{n}_2$ be space-like in $\mathbb{M}^5$. Then let

$$H_i = \{ \vec{v} \in \mathbb{M}^5 \mid (\vec{v}, \vec{n}_i) = 0 \}.$$

If $H_1 \cap H_2 \cap \mathbb{H}^4 \neq \emptyset$ and H_1 and H_2 intersect at angle θ , then

$$\frac{(\vec{n}_1, \vec{n}_2)}{\sqrt{(\vec{n}_1, \vec{n}_1)(\vec{n}_2, \vec{n}_2)}} = -\cos(\theta).$$

In particular, we will consider only examples where θ is always $\pi/2$, in which case the above simplifies to

$$(\vec{n}_1, \vec{n}_2) = 0.$$

Choose a representation $\rho \in \text{Hom}(\Gamma, G_4)$ sending each generator r_i to a reflection. In the above manner, we replace every reflection matrix $\rho(r_i)$ by a corresponding normal vector $\vec{n}_i$. We now have M Minkowski 5-vectors, instead of M 5×5 matrices. The relations of Γ written in terms of normal vectors all have the form:

$$(\vec{n}_i, \vec{n}_j) = 0 \text{ when } m_{ij} = 2 \text{ (angle condition)}$$

Let $U \subset \text{Hom}(\Gamma, G_4)$ be the open neighborhood of the inclusion map of Corollary 2.4. It is clear that, restricted to $\mu^{-1}(U)$, μ is a submersion with fibers given by scaling the normal vectors $\vec{n}_i$. Given a vector $v = \left(\dot{\vec{n}}_i \right)$ tangent to the point $(\vec{n}_i) \in \mu^{-1}(\rho)$, $\mu_* v \in T_\rho \mathbb{R}^{25M}$ is an infinitesimal deformation of ρ if and only if v satisfies the following derivatives of the above angle conditions:

$$\left(\dot{\vec{n}}_i, \vec{n}_j \right) + \left(\vec{n}_i, \dot{\vec{n}}_j \right) = 0 \text{ when } m_{ij} = 2$$

Any solution to this system of polynomials will now be a vector $(\dot{\vec{n}}_i)$ in $\mathbb{R}^{5M}$ instead of $\mathbb{R}^{25M}$. Note that we do not need to include the relation corresponding to each generator being an involution. This interpretation clearly has an immense computational advantage over working with the matrices directly.

Finally, to show ρ is infinitesimally rigid in $\text{Hom}(\Gamma, G_4)$ it suffices to show that the subspace of vectors $(\dot{\vec{n}}_i) \in \mathbb{R}^{5M}$ satisfying the above equations has dimension equal to $10 + M$, where $10 = \dim(G_4)$ dimensions come from infinitesimal conjugations by G_4 , and there are M dimensions corresponding to scaling the normal vectors, one for each generator of Γ .

3. THE 120-CELL

Here we will outline some of the basic properties of the hyperbolic 120-cell. We will use the word face to indicate a 2-cell, and wall to indicate a 3-cell. The 120-cell is a regular convex polytope formed by 120 dodecahedral walls, where we define regular to mean that its symmetry group acts transitively on the set of flags. (A flag of the 120-cell is quadruple consisting of 1 point, 1 edge containing the point, 1 pentagonal face containing the edge, and 1 dodecahedral wall containing the face.) The 120-cell has 600 vertices, and is dual to the 600-cell formed by 600 icosahedra.

The 120-cell can be embedded into $\mathbb{H}^4$ in such a way that the resulting convex hyperbolic polytope is compact, regular, and all intersecting dodecahedra hit at right angles. The quickest way to describe this polytope is by giving a list of 120 space-like vectors that are Minkowski-normal to the 120 walls. For completeness, we list the 120 normal vectors in $\mathbb{M}^5$, using the Golden ratio $\tau = (1 + \sqrt{5})/2$: [2]

- (1) The 8 vectors obtained by permuting the last 4 coordinates of $(\sqrt{2\tau}, \pm 2, 0, 0, 0)$.
- (2) The 16 vectors of the form $(\sqrt{2\tau}, \pm 1, \pm 1, \pm 1, \pm 1)$.
- (3) The 96 even permutations in the last 4 coordinates of $(\sqrt{2\tau}, \pm \tau, \pm 1, \pm \tau^{-1}, 0)$.

Consider the set $\mathcal{C}$ of walls of the 120-cell given by the 24 space-like vectors of items (1) and (2) above. Interestingly, these 24 vectors determine 24 hyperplanes of $\mathbb{H}^4$ which bound a regular (hyperideal) hyperbolic 24-cell, which is a polytope with 24 octahedral walls. This set $\mathcal{C}$ will play an important role here. One can compute directly that the walls of $\mathcal{C}$ are pairwise disjoint in $\mathbb{H}^4$. Moreover, there does not exist a set of 25 pairwise disjoint walls of the 120 cell.

Proposition 3.1. *Let P be the hyperbolic 120-cell. A maximum set of pairwise disjoint walls of P has 24 elements.*

Proof: Suppose we remove 24 walls from the 120-cell. For each wall that we remove, we place a marker on any wall which was adjacent to the removed wall. Each wall of the 120-cell is adjacent to 12 other walls; therefore after removing 24 walls we have placed $24 \times 12 = 288$ markers on the remaining 96 walls, or an average of 3 markers per wall that remains. The claim is that every remaining wall has exactly 3 markers. If this is indeed true, then removing an additional 25th wall would be impossible because any wall we removed would be adjacent to 3 of the walls in the original set of 24.

Suppose one of the walls had only 2 markers. This implies that there is at least one other wall with 4 markers, implying that the intersection of a certain set of 4 pairwise disjoint walls with this wall of 4 markers is a set of 4 pairwise disjoint pentagonal faces of a dodecahedron. This is impossible because the maximum number of pairwise disjoint faces of a dodecahedron is 3. Therefore no wall has 2 or fewer markers implying every wall has 3. $\square$

The above proof was explained to us by Allcock.

4. THE MAIN THEOREM

We are now ready to discuss the main result.

Theorem 4.1. *Let $P \subset \mathbb{H}^4$ be the hyperbolic regular right-angled 120-cell. Let $\mathcal{C}$ be the set of 24 pairwise disjoint walls of P given in Section 3. Let $\Gamma < G_4 =$*

$Isom(\mathbb{H}^4) < O(1,4)$ be the discrete infinite covolume group generated by the reflections in the 96 walls of P that are not in $\mathcal{C}$. Then the inclusion map $\Gamma \rightarrow G_4$ is infinitesimally rigid in the representation variety $Hom(\Gamma, G_4)$.

Before beginning the proof let us discuss the connection to Conjecture 1.1.

Proposition 4.2. *Let $\Gamma < G_4$ be the discrete group of Theorem 4.1. Then Γ has Fuchsian ends and the quotient C_Γ/Γ is isometric to the 120-cell P . In particular, Γ is convex cocompact.*

Proof: Consider the collection of 24 dodecahedral walls $\mathcal{C} \subset \mathbb{H}^4$ and its orbit $\Gamma \cdot \mathcal{C}$. Using the facts that all the dihedral angles of P are $\pi/2$, and all the walls of $\mathcal{C}$ are pairwise disjoint, it follows that any intersecting translates of $\mathcal{C}$ in the orbit $\Gamma \cdot \mathcal{C}$ glue together smoothly. In particular, the entire orbit is a disjoint union of totally geodesic hyperplanes. Let $C_\Gamma \subset \mathbb{H}^4$ be the Γ -invariant convex subset bounded by $\Gamma \cdot \mathcal{C}$. This shows that Γ has Fuchsian ends. It is clear that the quotient C_Γ/Γ is isometric to P . $\square$

Combining Theorem 4.1 and Proposition 4.2 shows that Conjecture 1.1 is true for Γ , which we record here.

Corollary 4.3. *The discrete subgroup $\Gamma < G_4$ has Fuchsian ends, is convex cocompact, and infinitesimally rigid, as predicted by Conjecture 1.1.*

Let us begin the proof of Theorem 4.1. We begin with 96 5×5 reflection matrices in $G_4 < O(1,4)$, the image of the generators of Γ under the inclusion map. We can replace each such reflection matrix with one of its (space-like) eigenvectors $\vec{n}_i$ corresponding to eigenvalue -1 . (All the possible choices are colinear.) The complete list of 96 vectors $\vec{n}_i$ will be the 96 even permutations in the last 4 coordinates of $(\sqrt{2}\tau, \pm\tau, \pm 1, \pm\tau^{-1}, 0)$, where τ is the Golden ratio. Pairs $\vec{n}_i$ and $\vec{n}_j$ corresponding to orthogonal walls of P will satisfy the relation

$$(\vec{n}_i, \vec{n}_j) = 0.$$

Let $W \subset T_{\vec{n}}\mathbb{R}^{480}$ be the linear space of infinitesimal deformations. Suppose we are given an infinitesimal deformation

$$\left(\dot{\vec{n}}_i\right) \in W \subset T_{\vec{n}}\mathbb{R}^{480} \cong \mathbb{R}^{480}$$

of the 96 space-like vectors which maps via μ_* to an infinitesimal deformation of Γ in G_4 . Then the following linear equation must hold for all pairs i, j corresponding to intersecting orthogonal walls of P :

$$\left(\dot{\vec{n}}_i, \vec{n}_j\right) + \left(\vec{n}_i, \dot{\vec{n}}_j\right) = 0.$$

Keep in mind that the 24 walls of $\mathcal{C}$ have been removed from P . There are 432 equations of this type, corresponding to the 432 remaining faces of P . As described above, we wish to find the solution space W of this set of linear polynomials. To prove that Γ is infinitesimally rigid it suffices to show that the solution space has $10 + 96$ dimensions, where 10 comes from the action of G_4 by conjugation, and 96 comes from scaling each of the 96 space-like vectors $\vec{n}_i$. These 96 scaling dimensions are killed by μ_* . Indeed, by this count we see that the solution space W has at least 106 dimensions, so it remains to show that it has at most 106 dimensions.

Begin by enumerating $\{p_1, p_2, \dots, p_{432}\}$ the polynomials defining W . Each p_i is a polynomial in the variables $(v_1, v_2, \dots, v_{480})$ where

$$\vec{n}_i = (v_{5(i-1)+1}, v_{5(i-1)+2}, v_{5(i-1)+3}, v_{5(i-1)+4}, v_{5i}).$$

Then define the matrix A_{alg} of algebraic numbers

$$(A_{\text{alg}})_{ij} := \frac{d}{dv_j} (p_i(v_1, v_2, \dots, v_{480})).$$

Each entry $(A_{\text{alg}})_{ij}$ is a number because each polynomial p_i is linear in the variables v_j . A_{alg} has 432 rows and 480 columns. The goal is to show its kernel has dimension at most 106. We will do this by showing the rank is at least 374.

In order to ensure that Mathematica can correctly and efficiently run the computation, we replace every algebraic number entry of A_{alg} with a square matrix of rational numbers. The number field $\mathbb{Q}(\alpha)$ contains every entry of A_{alg} when $\alpha = \sqrt{1 + \sqrt{5}}$. We choose the basis

$$\{1, \alpha, \alpha^2, \alpha^3\}$$

for $\mathbb{Q}(\alpha)$ as a vector space over $\mathbb{Q}$. Then each entry of A_{alg} is replaced by a 4×4 rational matrix representing its action by left multiplication on $\mathbb{Q}(\alpha)$ with respect to this basis. For example, the number $\frac{1}{2}(-1 - \sqrt{5})$ is replaced by the 4×4 matrix

$$\begin{pmatrix} 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & -2 \\ -\frac{1}{2} & 0 & -1 & 0 \\ 0 & -\frac{1}{2} & 0 & -1 \end{pmatrix}$$

Once we have the rational 4×4 matrix replacements for each entry of A_{alg} , it becomes a 432×480 matrix of 4×4 matrices. By ignoring the structure of the 4×4 matrices, we can think of it as a 1728×1920 matrix $A_{\mathbb{Q}}$ of rational numbers. It is easy to see that the kernel of $A_{\mathbb{Q}}$ has dimension at least 4 times that of A_{alg} . (If u is in the kernel of A_{alg} then $\{u, \alpha u, \alpha^2 u, \alpha^3 u\}$ will be in the kernel of $A_{\mathbb{Q}}$ and linearly independent over $\mathbb{Q}$.) Therefore the rank of $A_{\mathbb{Q}}$ is at most 4 times the rank of A_{alg} . To prove the infinitesimal rigidity of Γ , it therefore suffices to show that the rank of $A_{\mathbb{Q}}$ is at least $4 \cdot 374 = 1496$.

In order to further simplify the computation for Mathematica, multiply each row of $A_{\mathbb{Q}}$ by the least common multiple of the denominators of the rational numbers in that row. This gives us the integer matrix $A_{\mathbb{Z}}$ with the same rank as $A_{\mathbb{Q}}$. With some effort on a powerful computer (in 2008), Mathematica was able to determine that the rank of this matrix is 1496, the desired result. Therefore A_{alg} has a kernel of dimension at most 106. This proves that the inclusion map of the discrete group $\Gamma < G_4$ is infinitesimally rigid in the representation variety $\text{Hom}(\Gamma, G_4)$.

The rank calculation for $A_{\mathbb{Z}}$ took approximately 8 hours, and several gigabytes of RAM. This is perhaps related to the well known computational difficulty that diagonalizing an integer matrix via the Euclidean algorithm can lead to intermediate matrix entries of enormous size. We can greatly decrease the calculation time by reducing $A_{\mathbb{Z}}$ modulo a suitable prime p , obtaining matrix A_p . When working in the finite field $\mathbb{F}_p$ the diagonalization process is nearly trivial, and $\text{rank}(A_p) \leq \text{rank}(A_{\mathbb{Z}})$. Therefore, we simply reduce $A_{\mathbb{Z}}$ modulo several primes until we find one such that $\text{rank}(A_p) = 1496$. Once we find such a prime we are done by the above

inequality. Using the prime 113, this method allows us to arrive at the correct answer of a rank of 1496 for $A_{\mathbb{Z}}$ with a calculation time of approximately 5 seconds.

Acknowledgements: All computer calculations were performed in 2008 using Wolfram's Mathematica 6.0.1.0 on a MacPro running 64-bit Linux.

REFERENCES

- [1] E. Calabi. On compact Riemannian manifolds with constant curvature, I. In *Proc. Sympos. Pure Math., Vol. III*, pages 155–180. American Mathematical Society, Providence, R.I., 1961.
- [2] H.S.M. Coxeter. *Introduction to geometry*. Wiley Classics Library. John Wiley & Sons Inc., New York, 1989. Reprint of the 1969 edition.
- [3] H. Garland. On deformations of discrete groups in the noncompact case. In *Algebraic Groups and Discontinuous Subgroups (Proc. Sympos. Pure Math., Boulder, Colo., 1965)*, pages 405–412. Amer. Math. Soc., Providence, R.I., 1966.
- [4] W. Goldman. The symplectic nature of fundamental groups of surfaces. *Adv. in Math.*, (54):200–225, 1984.
- [5] S. Kerckhoff and P. Storm. From the 24-cell to the cuboctahedron. Preprint available at <http://front.math.ucdavis.edu/0805.4537>, 2008.
- [6] J. Ratcliffe. Hyperbolic manifolds. In *Handbook of geometric topology*, pages 899–920. North-Holland, Amsterdam, 2002.
- [7] W. Thurston. The topology and geometry of 3-manifolds. Available from the MSRI website www.msri.org, 1976–1979. Princeton Univ. lecture notes.
- [8] E.B. Vinberg and O.V. Shvartsman. Discrete groups of motions of spaces of constant curvature. In *Geometry. II.*, volume 29 of *Encyclopædia of Mathematical Sciences*, pages 139–248. Springer, Berlin, 1993.
- [9] A. Weil. On discrete subgroups of Lie groups. II. *Ann. of Math. (2)*, 75:578–602, 1962.